

für Zoonosen

The invasive vector mosquito Aedes japonicus in Europe: Still no end in sight!



Stephanie Thomas¹, Nils Tjaden¹, Anja Jaeschke^{1,2} Helge Kampen³, Doreen Werner⁴, Jürgen Kreyling¹, Carl Beierkuhnlein¹

¹Biogeography, University of Bayreuth, ²Biogeographical Modelling, University of Bayreuth, ³Friedrich-Loeffler-Institut, Federal Research Institute for Animal Health,


Greifswald – Insel Riems ⁴Leibniz Centre for Agricultural Landscape Research, Muencheberg; all Germany

Background

Aedes (Finlaya) japonicus japonicus (Theobald, 1901) is a highly ranked listed invasive culicid species and originates from East Asia and the Far East. At present it is established in the European countries Switzerland, Germany, Austria, Slovenia, Croatia, France, Belgium and the Netherlands as well as in America throughout the northeastern states of the USA crossing the border to the south of Canada, Hawaii, Washington and Oregon.


It is known to transmit the West Nile virus and Japanese Encephalitis virus under field conditions and is a potential vector for Dengue, Chikungunya, LaCrosse, Eastern Equine Encephalitis, St. Louis Encephalitis, and Rift Valley Virus.

Results: Areas at risk for an further establishment of Aedes japonicus




Climatic Variables	Explanation
Bio6	Min Temperature of Coldest Month
Bio9	Mean Temperature of Driest Quarter
Bio10	Mean Temperature of Warmest Quarter
Bio14	Precipitation of Warmest Month
Bio18	Precipitation of Warmest Quarter

Aedes japonicus - future




Large parts of Central and Eastern Europe will be climatically suitable

Haddow




depending on the ambient temperature.


Long-term establishments in Northern Europe and
Russia can be excluded.



The combination of correlative species distribution models with special expert knowledge on the ecology of a species, here an invasive diesease vector mosquito, can strongly improve the risk assessment regarding future areas at risk for an establishment of the species.

Climate Chamber Experiments: Egg Survival at Low Temperatures





Methods

- Correlative species distribution model SDMs: Ensemble Modelling approach as implemented in biomod2 R-package
- Bioclimatic variables of Worldclim (5' resolution, global scale), Generalized Linear Model (GLM), Generalized Boosting Model (GBM), Random Forest (RF),
 Maximum Entropy (MAXENT)
- Future projections 2030, 2050 and 2070 for five climate models (cesm1_bgc, fio_esm, giss_e2_r, inm_cm4, mpi_esm_lr) and two IPCC-5 scenarios (rcp4.5, rcp8.5)
- Climate chamber experiment: Detection of low temperature thresholds for mosquito egg survival under three different diurnal temperature range treatments and Combination of the outcome with SDMs to substantially improve risk maps by considering areas where an establishment of *Aedes japonicus* is unlikely due to winter conditions
- See also the talk of Anja Jaeschke SAT 14:30 CT2.7, H 21, RW II and the poster of Nils Tjaden P2.50

Further References

- Thomas, Beierkuhnlein (2013) Predicting ectotherm disease vector spread—benefits from multidisciplinary approaches and directions forward.

 Naturwissenschaften 100:395-405
- Kampen, Werner (2014) Out of the bush: the Asian bush mosquito Aedes japonicus japonicus (Theobald, 1901) (Diptera, Culicidae) becomes invasive.
 Parasites & Vectors 7:59
- Thomas, Obermayr et al. (2012) Low-temperature threshold for egg survival of a post-diapuase and non-diapause European aedine strain, Aedes
 albopictus (Diptera: Culicidae). Parasites & Vectors 5:100
- Fischer, Thomas et al. (2011) Projection of climatic suitability for Aedes albopictus Skuse (Culicidae) in Europe under climate change conditions. **Global Planetary Change** 78:54-64

Stephanie.Thomas@uni-bayreuth.de





www.biogeo.uni-bayreuth.de www.researchgate.net/profile/Stephanie_Thomas4